


Properties of Gases: Gas Pressure

- Gas Pressure

• Liquid Pressure
$$P \text{ (Pa)} = \frac{\text{Force (N)}}{\text{Area (m}^2\text{)}}$$

$$P = g \cdot h \cdot d$$


Barometric Pressure



Standard Atmospheric Pressure
1.00 atm
760 mm Hg, 760 torr
101.325 kPa
1.01325 bar
1013.25 mbar


The standard atmosphere is defined as the pressure exerted by a mercury column of exactly 760 mm in height when the density equals 13.6 g/cc.

Manometers



Simple Gas Laws


- Boyle 1662



Charles's Law

To convert Celsius to Kelvin:

$$T(\text{K}) = t(^{\circ}\text{C}) + 273.15$$



STP

- Gas properties depend on conditions.
- Standard conditions of temperature and pressure (STP).

$$P = 1 \text{ atm} = 760 \text{ mm Hg}$$

$$T = 0^\circ\text{C} = 273.15 \text{ K}$$

Avogadro's Law

- Equal volumes of gases at same temperature and pressure have equal numbers of molecules.
- Gas molecules may break up when they react.

Avogadro's Law

At a fixed temperature and pressure:

$$V \propto n \quad \text{or} \quad V = c n$$

At STP

$$1 \text{ mol gas} = 22.4 \text{ L gas}$$

Combining the Gas Laws: The Ideal Gas Equation and the General Gas Equation

- Boyle's law $V \propto 1/P$
 - Charles's law $V \propto T$
 - Avogadro's law $V \propto n$
- $$\left. \begin{array}{l} \\ \\ \end{array} \right\} V \propto \frac{nT}{P}$$

$$PV = nRT$$

The Gas Constant

$$PV = nRT$$

$$R = \frac{PV}{nT}$$

$$= 0.082057 \text{ L atm mol}^{-1} \text{ K}^{-1}$$

$$= 8.3145 \text{ m}^3 \text{ Pa mol}^{-1} \text{ K}^{-1}$$

$$= 8.3145 \text{ J mol}^{-1} \text{ K}^{-1}$$

The General Gas Equation

$$R = \frac{P_1 V_1}{n_1 T_1} = \frac{P_2 V_2}{n_2 T_2}$$

If we hold the number of moles and volume constant:

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

Molar Mass Determination

$$PV = nRT \quad \text{and} \quad n = \frac{m}{M}$$

$$PV = \frac{m}{M} RT$$


$$M = \frac{m RT}{PV}$$

$$M = \frac{dVRT}{PV}$$

Example

Using the Ideal gas Equation in Reaction Stoichiometry Calculations.

The decomposition of sodium azide, NaN_3 , at high temperatures produces $\text{N}_2(\text{g})$. Together with the necessary devices to initiate the reaction and trap the sodium metal formed, this reaction is used in air-bag safety systems. What volume of $\text{N}_2(\text{g})$, measured at 735 mm Hg and 26°C, is produced when 70.0 g NaN_3 is decomposed.



Example

Determine moles of N_2 :

$$n_{\text{N}_2} = 70 \text{ g N}_3 \times \frac{1 \text{ mol NaN}_3}{65.01 \text{ g N}_3/\text{mol N}_3} \times \frac{3 \text{ mol N}_2}{2 \text{ mol NaN}_3} = 1.62 \text{ mol N}_2$$

Determine volume of N_2 :


$$V = \frac{nRT}{P} = \frac{(1.62 \text{ mol})(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(299 \text{ K})}{\left\{ (735 \text{ mm Hg}) \frac{1.00 \text{ atm}}{760 \text{ mm Hg}} \right\}}$$
$$= 41.1 \text{ L}$$

Problem

If all gases are measured at the same temperature and pressure, what volume of $\text{NH}_3(\text{g})$ is produced when 225 L $\text{H}_2(\text{g})$ is consumed in the reaction $\text{N}_2(\text{g}) + \text{H}_2(\text{g}) \rightarrow \text{NH}_3(\text{g})$?

Dalton's Law of Partial Pressure

It states that the total pressure of a mixture of gases is the sum of the partial pressures of the components of the mixture.



Partial Pressure

$$P_{\text{tot}} = P_a + P_b + \dots$$

$$V_a = n_a RT / P_{\text{tot}} \quad \text{and} \quad V_{\text{tot}} = V_a + V_b + \dots$$

$$\frac{V_a}{V_{\text{tot}}} = \frac{n_a RT / P_{\text{tot}}}{n_{\text{tot}} RT / P_{\text{tot}}} = \frac{n_a}{n_{\text{tot}}}$$

$$\frac{P_a}{P_{\text{tot}}} = \frac{n_a RT / V_{\text{tot}}}{n_{\text{tot}} RT / V_{\text{tot}}} = \frac{n_a}{n_{\text{tot}}}$$

$$P_a = \chi_a \cdot P_{\text{tot}} \quad P_b = \chi_b \cdot P_{\text{tot}}$$


$$\frac{n_a}{n_{\text{tot}}} = \chi_a$$

Mole fraction of component A

Problem

The percent composition of air by volume is 78.08% N₂, 20.95% O₂, 0.93% Ar and 0.036% CO₂. What are the partial pressures of these four gases in a sample of air at a barometric pressure of 748 mm Hg?


Pneumatic Trough



$$P_{\text{tot}} = P_{\text{bar}} = P_{\text{gas}} + P_{\text{H}_2\text{O}}$$

Kinetic Molecular Theory


- Particles are point masses in constant, random, straight line motion.
- Particles are separated by great distances.
- Collisions are rapid and elastic.
- No force between particles.
- Total energy remains constant.



Average kinetic energy is directly proportional to temperature!!

Gas Properties Relating to the Kinetic-Molecular Theory

- Diffusion
 - Migration of molecules that results in a homogenous mixture.
- Effusion
 - Escape of gas molecules through a tiny pinhole.




Graham's Law

$$\frac{\text{rate of effusion of } A}{\text{rate of effusion of } B} = \frac{(u_{\text{rms}})_A}{(u_{\text{rms}})_B} = \sqrt{\frac{3RT/M_A}{3RT/M_B}} = \sqrt{\frac{M_B}{M_A}}$$


- Only for gases at low pressure (natural escape, not a jet).
- Tiny orifice (no collisions)
- Does not apply to diffusion.
 - Ratio used can be:
 - Rate of effusion (as above) – Distances traveled by molecules
 - Molecular speeds
 - Effusion times
 - Amounts of gas effused.

Real Gases

- Compressibility factor $PV/nRT = 1$
- Deviations occur for real gases.
 - $PV/nRT > 1$ - molecular volume is significant.
 - $PV/nRT < 1$ – intermolecular forces of attraction.



Real Gases



van der Waals Equation

$$\left(P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT$$
